

Microbiological examinations

Culture Media for use in the food industry, pharma
and cosmetics or veterinary medicine

Culture media at first-hand

FOR USE IN THE FOOD INDUSTRY,
PHARMA AND COSMETICS OR
VETERINARY MEDICINE



Culture media at first-hand

We are manufacturer of more than 200 different culture media and supplements to cultivate the microorganisms or specifically promote the growth of particularly fastidious species.

Thus, you are optimally equipped for the analysis of veterinary or human samples. Special media for food, water, pharmaceutical products complete the portfolio.

Do you have a specific request? Or are you looking for a supplier of media based on your own formulation?

We develop and produce according to your demand.

Our portfolio comprises:

- Dehydrated culture media
- Supplements and additives
- Ready-to-use culture media
- Culture media base materials
- Detection reagents

sifin diagnostics gmbh

Berliner Allee 317-321
13088 Berlin, Germany

Phone: +49 30 700 144-0
Telefax: +49 30 700 144-30
E-Mail: info@sifin.de

www.sifin.de

Contents

Miscellaneous

| | |
|--|----|
| Contact | 4 |
| Order blank | 6 |
| Certificate DIN EN ISO/IEC 17025 | 45 |

Culture media

| | |
|--|---------------|
| Culture media for food | 7 |
| Examination of milk, milk products and ice-cream | 8 |
| Examination of natural mineral, spring and table water ... | 13 |
| Water and sewage examination | 16 |
| Examination of drinking water | 19 |
| Examination of food | 21 |
| Examination of meat and meat products | 26 |
| Culture media for use in the beverage industry | 30 |
| Detection of anaerobic microorganisms | 32 |
| Detection of yeasts and moulds | 33 |
| Biochemical identification | 34 |
| Culture media for pharma | 35 |
| Culture media for cosmetics | 39 |
| Culture media for veterinary | 41 |

Your contact with sifin

You may reach us personally German time

Monday - Thursday

7:30 am to 4:15 pm

Friday

7:30 am to 3:00 pm

Head office

Telephone: +49 30 700 144-0

Telefax: +49 30 700 144-30

Export department

Telephone: +49 30 700 144-235 or
+49 30 700 144-283

Telefax: +49 30 700 144-30

E-Mail: info@sifin.de

Culture media

MICROBIOLOGY



Whether clinical diagnostics or industrial product control, we provide more than 200 different culture media and supplements to culture the microorganisms you are looking for, or to specifically promote the growth of particularly fastidious species. Thus, you are optimally equipped for analysis of food, water, pharmaceutical products, cosmetics as well as veterinary samples.

Culture media for different applications

Medical microbiology

- Culture media for growth and viable count
- Selective culture media
- Culture media for differentiation
- Culture media for sensibility testing

Microbiological examination of drinking water

- Culture media for colony count
- Media for the detection of:
Escherichia coli,
Enterococci,
Clostridium perfringens,
Pseudomonas aeruginosa

Culture media for the microbiological examinations in pharmaceutical and cosmetics industry

- Culture media for enrichment and growth
- Culture media for sterility testing
- Culture media for testing nonsterile products. Culture media are based on the recommendations of the harmonised method acc. to EP/USP/JP.

Special indications

- Environment
- Hygiene
- Fermentation

Fungi diagnostics

- Fungi, yeasts, dermatophytes
- Food and other test materials
- Pharmaceutical products acc. to harm. EP/USP/JP

Chromogenic culture media

- Dehydrated culture media

Veterinary medical microbiology

- Encourage development of fimbrial antigen F5 (K 99) in cattles: cultivation on Minca Agar and Minca Supplement
- Encourage development of fimbrial antigen F4 (K 88) in piglets: cultivation e. g. on Columbia Blood Agar with addition of sheep blood

Order Blank

sifin

Telefax +49 30 700 144-30

sifin diagnostics gmbh

Berliner Allee 317-321
13088 Berlin, Germany

Phone: +49 30 700 144-0
Telefax: +49 30 700 144-30
E-Mail: info@sifin.de

Customer no. _____

Name _____

| Art. No. | Quantity | Product |
|----------|----------|---------|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

Invoice address

Company / Department _____

Contact _____

Address _____

Zip, city / State _____

Telephone _____

Telefax _____

E-Mail _____

Delivery address

Company / Department _____

Contact _____

Address _____

Zip, city / State _____

Telephone _____

Telefax _____

E-Mail _____

☒ I order the items as mentioned above.

Date _____

Signature _____



Culture Media for Food

MICROBIOLOGY



Trust is fine, control is best. Culture media for food examinations.

Contents

- Examination of milk, milk products and ice-cream
- Examination of natural mineral, spring and table water
- Water and sewage examination
- Examination of drinking water
- Examination of food
- Examination of meat and meat products
- Culture media for use in the beverage industry
- Detection of anaerobic microorganisms
- Detection of yeasts and moulds
- Biochemical identification



Examination of Milk, Milk Products and Ice-Cream

E. coli and coliforms

enrichment

| Art. No. | Product | Description | Packing |
|----------|--|---|---------|
| TN1109 | Brilliant Green Bile Lactose Broth (BRILA Broth) | for 12.5 l culture medium: single strength (40.0 g for 1 l medium) for 6.3 l culture medium: double strength (80.0 g for 1 l medium) | 500 g |
| TN1152 | Lauryl Sulfate Broth (Lauryl Sulfate Tryptose Lactose Broth) | for 14.0 l culture medium (35.6 g for 1 l culture medium) | 500 g |
| TN1153 | Lauryl Sulfate Tryptose Broth with Tryptophan and MU (LST/MUG Medium) | for 13.6 l culture medium: single strength (36.7 g for 1 l medium) for 6.8 l culture medium: double strength (73.4 g for 1 l medium) | 500 g |

isolation

| | | | |
|---------------------|---|---|---------------|
| TN1130 | Endo Agar (Lactose Fuchsin Sulfite Agar) | for 14.5 l culture medium (34.6 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1156 | MacConkey Agar | for 10.3 l culture medium (48.5 g for 1 l culture medium) | 500 g |
| TN1075 TN1075-01 | MacConkey Agar acc. to harm. EP/USP/JP | for 10.0 l culture medium (50.0 g for 1 l culture medium) for 100.0 l culture medium (50.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1259 | TBX Chromogen Agar (Tryptone Bile X Glucuronic Agar) | for 15.8 l culture medium (31.6 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1042 | Violet Red Bile Glucose Agar acc. to harm. EP/USP/JP (VRBD Agar) | for 13.7 l culture medium (36.5 g for 1 l culture medium) | 500 g |
| TN1149 | Violet Red Bile Lactose Agar (VRBL Agar, VRB Agar) | for 13.3 l culture medium (37.5 g for 1 l culture medium) | 500 g |
| TN1148 | VRBD Agar (VRBD Agar) | for 13.3 l culture medium (37.5 g for 1 l culture medium) | 500 g |

Salmonella

non-selective pre-enrichment

| Art. No. | Product | Description | Packing |
|---------------------|-------------------------------------|---|---------------|
| TN1137 TN1137-01 | Buffered Peptone Water (BPW) | for 25 l culture medium (20.0 g for 1 l culture medium) for 250 l culture medium (20.0 g for 1 l culture medium) | 500 g 5 kg |

selective enrichment

| | | | |
|--------|--|---|-------|
| TN1157 | Magnesium Chloride Malachite Green Broth acc. to Rappaport-Vassiliadis (RV Medium) | for 15.0 l culture medium (33.4 g for 1 l culture medium) | 500 g |
| TN1262 | Rappaport Vassiliadis Salmonella Enrichment Broth acc. to harm. EP/USP/JP (RVS Broth) | for 18.5 l culture medium (27.1 g for 1 l culture medium) | 500 g |
| TN1183 | Selenite Cystine Broth | for 21.7 l culture medium (23.0 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1187 | Tetrathionate Broth (Base) acc. to Muller-Kauffmann | for 5.0 l culture medium (89.4 g for 1 l culture medium) Addition of iodine-potassium iodide solution (5 g potassium iodide, 4 g iodine, 20 ml distilled water). | 500 g |

isolation

| | | | |
|---------------------|---|---|---------------|
| TN1110 | Brilliant Green Phenol Red Agar acc. to Edel Kampelmacher (BPLS Agar, mod.) | for 10.1 l culture medium (49.7 g for 1 l culture medium) | 500 g |
| TN1121 | Desoxycholate Citrate Agar Modified (Leifson Agar, DCLS Agar, mod.) | for 10.1 l culture medium (49.5 g for 1 l culture medium) | 500 g |
| TN1221 TN1221-01 | SS Agar (Salmonella Shigella Agar Modified) | for 8.8 l culture medium (56.5 g for 1 l culture medium) for 88.5 l culture medium (56.5 g for 1 l culture medium) | 500 g 5 kg |
| TN1194 TN1194-01 | Water-blue Metachrome-yellow Lactose Agar acc. to Gassner, Modified (Gassner Agar, mod.) | for 9.9 l culture medium (50.3 g for 1 l culture medium) for 99.4 l culture medium (50.3 g for 1 l culture medium) | 500 g 5 kg |
| TN1170 | XLD Agar acc. to harm. EP/USP/JP (Xylose Lysine Deoxycholate Agar) | for 9.1 l culture medium (55.2 g for 1 l culture medium) | 500 g |
| TN1196 | XLD Agar acc. to ISO 6579 (Xylose Lysine Deoxycholate Agar) | for 8.8 l culture medium (57.0 g for 1 l culture medium) | 500 g |

confirmation

| | | | |
|--------|--|--|-------------|
| TN1146 | Kligler Iron Agar (KIA, Double Sugar Iron Agar acc. to Kligler) | for 10.1 l culture medium (49.6 g for 1 l culture medium) Urea TN1308 may be added. | 500 g |
| TN1154 | Lysine Decarboxylase Broth (LDC Broth) | for 5.5 l culture medium (9.0 g for 1 l culture medium) | 50 g |
| TN1272 | MSRV Medium (Base) (Modified Semi-Solid Rappaport Vassiliadis Medium) | for 15.8 l culture medium (31.6 g for 1 l culture medium) Addition of Novobiocin Selective Supplement TN1324. | 500 g |
| TN1324 | Novobiocin Selective Supplement | for 6 l culture medium (2 vials for 1 l culture medium) 1 vial for 0.5 l MSRV Medium (Base) TN1272. | 12 x 1 vial |
| TN1308 | Urea Solution, 40 % | Supplement for culture medium: 1 vial for 1 l Kligler Iron Agar TN1146. | 12 x 1 vial |

Coagulase-positive staphylococci

| Art. No. | Product | Description | Packing |
|----------|---|---|-----------|
| TN1104 | Baird Parker Agar (Base) | for 7.9 l culture medium (63.0 g for 1 l culture medium) Addition of Egg Yolk Tellurite Emulsion TN1310. | 500 g |
| TN1310 | Egg Yolk Potassium Tellurite Emulsion | for approx. 2 l culture medium (2 vials for 1 l culture medium) Supplement for culture media: 1 vial contains 25 ml 2 vials for 950 ml Baird Parker Agar (Base) TN1104. | 4 x 25 ml |
| TN1166 | Mannitol Salt Agar acc. to harm. EP/USP/JP | for 4.5 l culture medium (111.0 g for 1 l culture medium) | 500 g |

Yeasts and moulds

| Art. No. | Product | Description | Packing |
|---------------------|--|---|---------------|
| TN1424 | Glycerin (water-free) | Supplement for culture media: Wort Agar (Base) TN1210 | 100 ml |
| TN1264 | Sabouraud Dextrose Agar acc. to harm. EP/USP/JP | for 7.7 l culture medium (65.0 g for 1 l culture medium) | 500 g |
| TN1178 TN1178-01 | Sabouraud 2% Glucose Agar (Sabouraud Dextrose Agar) | for 11.9 l culture medium (42.0 g for 1 l culture medium) for 119.0 l culture medium (42.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1746 | Wort Agar | 1 bottle contains 250 ml. | 250 ml |
| TN1210 | Wort Agar (Base) | for 10.4 l culture medium (48.0 g for 1 l culture medium) Addition of glycerin TN1424. | 500 g |
| TN1144 | Yeast Extract Glucose Chloramphenicol Agar (YGC Agar) | for 14.2 l culture medium (35.1 g for 1 l culture medium) No supplements necessary. | 500 g |

Acid and non-acid forming microorganisms

| Art. No. | Product | Description | Packing |
|----------|--------------------------------|---|---------|
| TN1114 | China Blue Lactose Agar | for 13.0 l culture medium (38.5 g for 1 l culture medium) | 500 g |

Clostrides

| Art. No. | Product | Description | Packing |
|------------------|---|---|---------------|
| TN1116 | Clostridial Differential Broth (DRCM) | for 16.6 l culture medium: single strength (30.1 g for 1 l medium) for 8.3 l culture medium: double strength (60.2 g for 1 l medium) | 500 g |
| TN1167 | Reinforced Medium for Clostridia acc. to harm. EP/USP/JP (RCM) | for 13.2 l culture medium (38.0 g for 1 l culture medium) | 500 g |
| TN1188 TN1253 | Thioglycollate Medium acc. to EP/USP (Fluid Thioglycollate Medium) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |

Listeria

| Art. No. | Product | Description | Packing |
|---------------------|--|--|---------------|
| TN1257 | Fraser Broth (Base) | for 8.7 l culture medium (57.4 g for 1 l culture medium) Addition of Fraser Selective Supplement TN1318 or ½ Fraser Selective Supplement TN1319. | 500 g |
| TN1035 | Fraser Broth Complete | for 8.6 l culture medium (57.9 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1034 TN1034-01 | ½ Fraser Broth Complete | for 8.6 l culture medium (57.9 g for 1 l culture medium) for 86.3 l culture medium (57.9 g for 1 l culture medium) No supplements necessary. | 500 g 5 kg |
| TN1318 | Fraser Selective Supplement | for 3 l culture medium (1 vial per component for 0.5 l medium) Supplement for culture media consists of two components. 1 vial per component for 0.5 l Fraser Broth (Base) TN1257. | 6 x 2 vials |
| TN1209 | Palcam Agar (Base) (Palcam Agar acc. to van Netten) | for 7.4 l culture medium (68.0 g for 1 l culture medium) Addition of Palcam Selective Supplement TN1312. | 500 g |
| TN1312 | Palcam Selektive Supplement | for 6 l culture medium (2 vials for 1 l culture medium) 1 vial for 0.5 l Palcam Agar (Base) TN1209. | 12 x 1 vial |

Bacillus cereus

| Art. No. | Product | Description | Packing |
|----------|--|--|-------------|
| TN1245 | Bacillus Cereus Agar (Base) (PEMBA) | for 12.5 l culture medium (39.9 g for 1 l culture medium) Addition of Egg Yolk Emulsion TN1316 and Bacillus Cereus Selective Supplement TN1315. | 500 g |
| TN1315 | Bacillus Cereus Selective Supplement (Polymyxin B Selective Supplement) | for 6 l culture medium (2 vials for 1 culture medium) In conjunction with Bacillus Cereus Agar (Base) TN1245 and Egg Yolk Emulsion TN1316. | 12 x 1 vial |
| TN1316 | Egg Yolk Emulsion | for approx. 1-2 l culture medium (2-4 vials for 1 l culture medium) Supplement for culture media: 1 vial contains 25 ml 2 vials for 950 ml Bacillus Cereus Agar (Base) TN1245. 4 vials for 900 ml MYP Agar (Base) TN1291. | 4 x 25 ml |
| TN1291 | MYP Agar (Base) (Cereus Selective Agar Base acc. to Mossel) | for 11.6 l culture medium (43.0 g for 1 l culture medium) Addition of Egg Yolk Emulsion TN1316 and Bacillus Cereus Selective Supplement TN1315. | 500 g |

Examination of Natural Mineral, Spring and Table Water

Germ number determination

| Art. No. | Product | Description | Packing |
|---------------------|--|--|---------------|
| TN1124 | DEV Gelatin Agar | for 11.1 l culture medium (45.0 g for 1 l culture medium) | 500 g |
| TN1724 | DEV Nutrient Agar | 1 bottle contains 250 ml. | 250 ml |
| TN1128 TN1128-01 | DEV Nutrient Agar | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1164 TN1164-01 | Nutrient Agar I | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) Blood may be added. | 500 g 5 kg |
| TN1176 | Nutrient Agar acc. to the German drinking water regulation and Food and Feed Code | 25.0 g for 1 l culture medium (25.0 g for 1 l culture medium) Blood may be added. | 500 g |
| TN1189 | Plate Count Agar (TGE Agar, Tryptone Glucose Yeast Extract Agar) | for 27.0 l culture medium (18.5 g for 1 l culture medium) | 500 g |

E. coli and coliform germs

enrichment

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1109 | Brilliant Green Bile Lactose Broth (BRILA Broth) | for 12.5 l culture medium: single strength (40.0 g for 1 l medium) for 6.3 l culture medium: double strength (80.0 g for 1 l medium) | 500 g |
| TN1127 TN1127-01 | DEV Lactose Peptone Broth | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1152 | Lauryl Sulfate Broth (Lauryl Sulfate Tryptose Lactose Broth) | for 14.0 l culture medium (35.6 g for 1 l culture medium) | 500 g |
| TN1153 | Lauryl Sulfate Tryptose Broth with Tryptophan and MUG (LST/MUG Medium) | for 13.6 l culture medium: single strength (36.7 g for 1 l medium) for 6.8 l culture medium: double strength (73.4 g for 1 l medium) | 500 g |

isolation

| | | | |
|---------------------|---|---|---------------|
| TN1130 | Endo Agar (Lactose Fuchsin Sulfite Agar) | for 14.5 l culture medium (34.6 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1156 | MacConkey Agar | for 10.3 l culture medium (48.5 g for 1 l culture medium) | 500 g |
| TN1075 TN1075-01 | MacConkey Agar acc. to harm. EP/USP/JP | for 10.0 l culture medium (50.0 g for 1 l culture medium) for 100.0 l culture medium (50.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1259 | TBX Chromogen Agar (Tryptone Bile X Glucuronic Agar) | for 15.8 l culture medium (31.6 g for 1 l culture medium) No supplements necessary. | 500 g |

confirmation

| Art. No. | Product | Description | Packing |
|----------|---|--|-------------|
| TN1517 | Oxidase reagent | for 12 tests 1 vial in 2 ml distilled water. | 12 x 1 vial |
| TN1258 | Tryptophan Pepton Water (Tryptophan Broth) | for 6.3 l culture medium (16.0 g for 1 l culture medium) | 100 g |

Faecal streptococci

enrichment

| Art. No. | Product | Description | Packing |
|----------|----------------------------|---|---------|
| TN1103 | Azide Glucose Broth | for 14.3 l culture medium: single strength (35.0 g for 1 l medium) for 7.4 l culture medium: double strength (70.0 g for 1 l medium) | 500 g |

isolation

| | | | |
|--------|--|--|-------|
| TN1132 | Enterococci Selective Agar acc. to Slanetz-Bartley (Enterococci Agar) | for 12.0 l culture medium (41.5 g for 1 l culture medium) | 500 g |
| TN1217 | Kanamycin Esculin Azide Agar (KAA Agar) | for 11.7 l culture medium (42.7 g for 1 l culture medium) No supplements necessary. | 500 g |

confirmation

| | | | |
|--------|--|--|-------|
| TN1276 | Bile Esculin Azide Agar (BEM, Bile Esculin Medium) | for 9.3 l culture medium (53.7 g for 1 l culture medium) | 500 g |
| TN1216 | Brain Heart Infusion Broth (BHI Broth, Brain Heart Glucose Broth) | for 13.5 l culture medium (37.0 g for 1 l medium) | 500 g |

Pseudomonas aeruginosa

enrichment

| Art. No. | Product | Description | Packing |
|----------|------------------------------|---|---------|
| TN1158 | Malachite-green Broth | for 56.8 l culture medium (8.8 g for 1 l culture medium) No supplements necessary. | 500 g |

isolation

| | | | |
|--------|--|--|-------------|
| TN1424 | Glycerin (water-free) | Supplement for culture media: Pseudomonas Agar (Base) TN1286 | 100 ml |
| TN1517 | Oxidase reagent | for 12 tests 1 vial in 2 ml distilled water. | 12 x 1 vial |
| TN1286 | Pseudomonas Agar (Base) | for 10.3 l culture medium (48.4 g for 1 l culture medium) Addition of Pseudomonas CN Selective Supplement TN1323 and glycerin TN1424. | 500 g |
| TN1323 | Pseudomonas CN Selective Supplement | for 6 l culture medium (2 vials for 1 l culture medium) 1 vial for 0.5 l Pseudomonas Agar (Base) TN1286. | 12 x 1 vial |

Sulphite-reducing, anaerobic spore formers

enrichment

| Art. No. | Product | Description | Packing |
|----------|---|---|---------|
| TN1116 | Clostridial Differential Broth (DRCM) | for 16.6 l culture medium: single strength (30.1 g for 1 l medium) for 8.3 l culture medium: double strength (60.2 g for 1 l medium) | 500 g |
| TN1167 | Reinforced Medium for Clostridia acc. to harm. EP/USP/JP (RCM) | for 13.2 l culture medium (38.0 g for 1 l culture medium) | 500 g |

confirmation

| | | | |
|---------------------|--|---|---------------|
| TN1288 | m-CP Agar (Base) (Chromogenic Membrane Filtration Clostridium perfringens Agar) | for 7.3 l culture medium (68.2 g for 1 l culture medium) Addition of m-CP-Selective Supplement TN1330. | 500 g |
| TN1330 | m-CP Selective Supplement | for 2 l culture medium (1 vial per component for 0.5 l culture medium) Supplement for culture media consists of three components. 1 vial per component for 0.5 l m-CP Agar (Base) TN1288. | 4 x 3 vials |
| TN1188 TN1188-01 | Thioglycollate Medium acc. to EP/USP (Fluid Thioglycollate Medium) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |

Water and Sewage Examination

Germ number determination

| Art. No. | Product | Description | Packing |
|---------------------|---|--|---------------|
| TN1124 | DEV Gelatin Agar | for 11.1 l culture medium (45.0 g for 1 l culture medium) | 500 g |
| TN1724 | DEV Nutrient Agar | 1 bottle contains 250 ml. | 250 ml |
| TN1128 TN1128-01 | DEV Nutrient Agar | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1164 TN1164-01 | Nutrient Agar I | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) Blood may be added. | 500 g 5 kg |
| TN1189 | Plate Count Agar (TGE Agar, Tryptone Glucose Yeast Extract Agar) | for 27.0 l culture medium (18.5 g for 1 l culture medium) | 500 g |

E. coli and coliforms

enrichment

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1109 | Brilliant Green Bile Lactose Broth (BRILA Broth) | for 12.5 l culture medium: single strength (40.0 g for 1 l medium) for 6.3 l culture medium: double strength (80.0 g for 1 l medium) | 500 g |
| TN1127 TN1127-01 | DEV Lactose Peptone Broth | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1152 | Lauryl Sulfate Broth (Lauryl Sulfate Tryptose Lactose Broth) | for 14.0 l culture medium (35.6 g for 1 l culture medium) | 500 g |
| TN1153 | Lauryl Sulfate Tryptose Broth with Tryptophan and MUG (LST/MUG Medium) | for 13.6 l culture medium: single strength (36.7 g for 1 l medium) for 6.8 l culture medium: double strength (73.4 g for 1 l medium) | 500 g |

isolation

| | | | |
|---------------------|---|---|---------------|
| TN1130 | Endo Agar (Lactose Fuchsin Sulfite Agar) | for 14.5 l culture medium (34.6 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1156 | MacConkey Agar | for 10.3 l culture medium (48.5 g for 1 l culture medium) | 500 g |
| TN1075 TN1075-01 | MacConkey Agar acc. to harm. EP/USP/JP | for 10.0 l culture medium (50.0 g for 1 l culture medium) for 100.0 l culture medium (50.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1042 | Violet Red Bile Glucose Agar acc. to harm. EP/USP/JP (VRBD Agar) | for 13.7 l culture medium (36.5 g for 1 l culture medium) | 500 g |
| TN1149 | VRB Agar (VRBL Agar, VRB Agar) | for 13.3 l culture medium (37.5 g for 1 l culture medium) | 500 g |
| TN1148 | VRBD Agar (Violet Red Bile Glucose Agar) | for 13.3 l culture medium (37.5 g for 1 l culture medium) | 500 g |

confirmation

| Art. No. | Product | Description | Packing |
|----------|---|--|-------------|
| TN1517 | Oxidase reagent | for 12 tests: 1 vial in 2 ml distilled water. | 12 x 1 vial |
| TN1258 | Tryptophan Pepton Water (Tryptophan Broth) | for 6.3 l culture medium (16.0 g for 1 l culture medium) | 100 g |

Faecal streptococci

enrichment

| Art. No. | Product | Description | Packing |
|----------|--|---|---------|
| TN1103 | Azide Glucose Broth | for 14.3 l culture medium: single strength (35.0 g for 1 l medium) for 7.4 l culture medium: double strength (70.0 g for 1 l medium) | 500 g |
| TN1218 | Kanamycin Esculin Azide Broth (KAA Broth) | for 15.3 l culture medium (32.7 g for 1 l culture medium) No supplements necessary. | 500 g |

isolation

| | | | |
|--------|--|--|-------|
| TN1132 | Enterococci Selective Agar acc. to Slanetz-Bartley (Enterococci Agar) | for 12.0 l culture medium (41.5 g for 1 l culture medium) | 500 g |
| TN1217 | Kanamycin Esculin Azide Agar (KAA Agar) | for 11.7 l culture medium (42.7 g for 1 l culture medium) No supplements necessary. | 500 g |

confirmation

| | | | |
|--------|---|--|-------|
| TN1276 | Bile Esculin Azide Agar (BEM, Bile Esculin Medium) | for 9.3 l culture medium (53.7 g for 1 l culture medium) | 500 g |
|--------|---|--|-------|

Pseudomonas aeruginosa

enrichment

| Art. No. | Product | Description | Packing |
|----------|------------------------------|---|---------|
| TN1158 | Malachite-green Broth | for 56.8 l culture medium (8.8 g for 1 l culture medium) No supplements necessary. | 500 g |

isolation

| | | | |
|--------|--|--|-------------|
| TN1424 | Glycerin (water-free) | Supplement for culture media: Pseudomonas Agar (Base) TN1286 | 100 ml |
| TN1517 | Oxidase reagent | for 12 tests 1 vial in 2 ml distilled water. | 12 x 1 vial |
| TN1286 | Pseudomonas Agar (Base) | for 10.3 l culture medium (48.4 g for 1 l culture medium) Addition of Pseudomonas CN Selective Supplement TN1323 and glycerin TN1424. | 500 g |
| TN1323 | Pseudomonas CN Selective Supplement | for 6 l culture medium (2 vials for 1 l culture medium) 1 vial for 0.5 l Pseudomonas Agar (Base) TN1286. | 12 x 1 vial |

Sulphite-reducing, anaerobic spore formers

enrichment

| Art. No. | Product | Description | Packing |
|----------|---|---|---------|
| TN1116 | Clostridial Differential Broth (DRCM) | for 16.6 l culture medium: single strength (30.1 g for 1 l medium) for 8.3 l culture medium: double strength (60.2 g for 1 l medium) | 500 g |
| TN1167 | Reinforced Medium for Clostridia acc. to harm. EP/USP/JP (RCM) | for 13.2 l culture medium (38.0 g for 1 l culture medium) | 500 g |

confirmation

| | | | |
|---------------------|--|---|---------------|
| TN1288 | m-CP Agar (Base) (Chromogenic Membrane Filtration Clostridium perfringens Agar) | for 7.3 l culture medium (68.2 g for 1 l culture medium) Addition of m-CP-Selective Supplement TN 1330. | 500 g |
| TN1188 TN1188-01 | Thioglycollate Medium acc. to EP/USP (Fluid Thioglycollate Medium) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |

Staphylococcus aureus

enrichment

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1263 TN1263-01 | Casein Soya Bean Digest Broth acc. to harm. EP/USP/JP (CASO Broth acc. to harm. EP/USP/JP) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |

isolation

| | | | |
|--------|---|---|-----------|
| TN1104 | Baird Parker Agar (Base) | for 7.9 l culture medium (63.0 g for 1 l culture medium) Addition of Egg Yolk Tellurite Emulsion TN1310. | 500 g |
| TN1310 | Egg Yolk Potassium Tellurite Emulsion | for approx. 2 l culture medium (2 vials for 1 l culture medium) Supplement for culture media: 1 vial contains 25 ml 2 vials for 950 ml Baird Parker Agar (Base) TN1104. | 4 x 25 ml |
| TN1166 | Mannitol Salt Agar acc. to harm. EP/USP/JP | for 4.5 l culture medium (111.0 g for 1 l culture medium) | 500 g |

Examination of Drinking Water

E. coli and coliforms

| Art. No. | Product | Description | Packing |
|---------------------|--|--|---------------|
| TN1064 | ChroMedium Coliform (Chromogenic Coliform Agar, CCA) | for 17.3 l culture medium (28.9 g for 1 l culture medium) | 500 g |
| TN1031 TN1031-01 | Casein Soya Bean Digest Agar acc. to harm. EP/USP/JP (CASO Agar) | for 12.5 l culture medium (40.0 g for 1 l culture medium) for 125.0 l culture medium (40.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1724 | DEV Nutrient Agar | 1 bottle contains 250 ml. | 250 ml |
| TN1128 TN1128-01 | DEV Nutrient Agar | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1278 | Lactose TTC Agar with Tergitol-7 (Tergitol-7 TTC Agar) | for 9.4 l culture medium (53.2 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1164 TN1164-01 | Nutrient Agar I | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) Blood may be added. | 500 g 5 kg |
| TN1176 | Nutrient Agar acc. to the German drinking water regulation and Food and Feed Code | for 20 l culture medium (25.0 g for 1 l culture medium) Blood may be added. | 500 g |
| TN1517 | Oxidase reagent | for 12 tests, 1 vial in 2 ml distilled water. | 12 x 1 vial |
| TN1258 | Tryptophan Pepton Water (Tryptophan Broth) | for 6.3 l culture medium (16.0 g for 1 l culture medium) | 100 g |

Enterococci

| Art. No. | Product | Description | Packing |
|----------|--|---|---------|
| TN1276 | Bile Esculin Azide Agar (BEM, Bile Esculin Medium) | for 9.3 l culture medium (53.7 g for 1 l culture medium) | 500 g |
| TN1132 | Enterococci Selective Agar acc. to Slanetz-Bartley (Enterococci Agar) | for 12.0 l culture medium (41.5 g for 1 l culture medium) | 500 g |

Clostrides

| Art. No. | Product | Description | Packing |
|----------|--|---|-----------------|
| TN1519 | Acid Phosphatase Reagent | 6 vials dye reagent, 1 bottle buffer | 6 vials + 1 btl |
| TN1288 | m-CP Agar (Base) (Chromogenic Membrane Filtration Clostridium perfringens Agar) | for 7.3 l culture medium (68.2 g for 1 l culture medium) Addition of m-CP-Selective Supplement TN1330. | 500 g |
| TN1330 | m-CP Selective Supplement | for 2 l culture medium (1 vial per component for 0.5 l culture medium) Supplement for culture media consists of three components. 1 vial per component for 0.5 l m-CP Agar (Base) TN1288. | 4 x 3 vials |
| TN1241 | TSC Agar (Base) (Tryptose Sulfite Cycloserine Agar) | for 12.8 l culture medium (39.0 g for 1 l culture medium) Addition of TSC Selective Supplement TN1333. | 500 g |
| TN1333 | TSC Selective Supplement | for 6 l culture medium (2 vials for 1 l culture medium) 1 vial for 0.5 l TSC Agar (Base) TN1241. | 12 x 1 vial |

Pseudomonads

| Art. No. | Product | Description | Packing |
|----------|--|--|-------------|
| TN1424 | Glycerin (water-free) | Supplement for culture media: King's Agar (Base) acc. to EN 12780 TN 1287 Pseudomonas Agar (Base) TN1286 | 100 ml |
| TN1287 | King's Agar (Base) | for 2.7 l culture medium (36.9 g for 1 l culture medium) Addition of glycerin TN1424. | 100 g |
| TN1176 | Nutrient Agar acc. to the German drinking water regulation and Food and Feed Code | for 20 l culture medium (25.0 g for 1 l culture medium) Blood may be added. | 500 g |
| TN1517 | Oxidase reagent | for 12 tests 1 vial in 2 ml distilled water. | 12 x 1 vial |
| TN1286 | Pseudomonas Agar (Base) | for 10.3 l culture medium (48.4 g for 1 l culture medium) Addition of Pseudomonas CN Selective Supplement TN1323 and glycerin TN1424. | 500 g |
| TN1323 | Pseudomonas CN Selective Supplement | for 6 l culture medium (2 vials for 1 l culture medium) 1 vial for 0.5 l Pseudomonas Agar (Base) TN1286. | 12 x 1 vial |

Total germ count

| Art. No. | Product | Description | Packing |
|---------------------|---------------------------|---|---------------|
| TN1724 | DEV Nutrient Agar | 1 bottle contains 250 ml. | 250 ml |
| TN1128 TN1128-01 | DEV Nutrient Agar | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1290 | Yeast Extract Agar | for 26.3 l culture medium (19.0 g for 1 l culture medium) | 500 g |

Examination of Food

Germ number determination

| Art. No. | Product | Description | Packing |
|---------------------|---|--|---------------|
| TN1164 TN1164-01 | Nutrient Agar I | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) Blood may be added. | 500 g 5 kg |
| TN1189 | Plate Count Agar (TGE Agar, Tryptone Glucose Yeast Extract Agar) | for 27.0 l culture medium (18.5 g for 1 l culture medium) | 500 g |

Enterobacteriaceae

pre-enrichment

| Art. No. | Product | Description | Packing |
|---------------------|-------------------------------------|---|---------------|
| TN1137 TN1137-01 | Buffered Peptone Water (BPW) | for 25.0 l culture medium (20.0 g for 1 l culture medium) for 250.0 l culture medium (20.0 g for 1 l culture medium) | 500 g 5 kg |

isolation

| | | | |
|---------------------|--|---|---------------|
| TN1042 | Violet Red Bile Glucose Agar acc. to harm. EP/USP/JP (VRBD Agar) | for 13.7 l culture medium (36.5 g for 1 l culture medium) | 500 g |
| TN1148 | VRBD Agar (Violet Red Bile Glucose Agar) | for 13.3 l culture medium (37.5 g for 1 l culture medium) | 500 g |
| TN1194 TN1194-01 | Water-blue Metachrome-yellow Lactose Agar acc. to Gassner Modified (Gassner Agar, mod.) | for 9.9 l culture medium (50.3 g for 1 l culture medium) for 99.4 l culture medium (50.3 g for 1 l culture medium) | 500 g 5 kg |

confirmation

| | | | |
|---------------------|--|---|---------------|
| TN1200 | Eosine Methylene Blue Lactose Saccharose Agar (EMB Agar, Levine Agar) | for 15.4 l culture medium (32.5 g for 1 l culture medium) | 500 g |
| TN1146 | Kligler Iron Agar (KIA, Double Sugar Iron Agar acc. to Kligler) | for 10.1 l culture medium (49.6 g for 1 l culture medium) Urea Solution, 40% TN 1308 may be added. | 500 g |
| TN1156 | MacConkey Agar | for 10.3 l culture medium (48.5 g for 1 l culture medium) | 500 g |
| TN1075 TN1075-01 | MacConkey Agar acc. to harm. EP/USP/JP | for 10.0 l culture medium (50.0 g for 1 l culture medium) for 100.0 l culture medium (50.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1184 | Simmons Citrate Agar | for 9.8 l culture medium (15.3 g for 1 l culture medium) | 150 g |

E. coli

enrichment

| Art. No. | Product | Description | Packing |
|----------|---|---|---------|
| TN1109 | Brilliant Green Bile Lactose Broth (BRILA Broth) | for 12.5 l culture medium: single strength (40.0 g for 1 l medium) for 6.3 l culture medium: double strength (80.0 g for 1 l medium) | 500 g |
| TN1152 | Lauryl Sulfate Broth (Lauryl Sulfate Tryptose Lactose Broth) | for 14.0 l culture medium (35.6 g for 1 l culture medium) | 500 g |
| TN1153 | Lauryl Sulfate Tryptose Broth with Tryptophan and MUG (LST/MUG Medium) | for 13.6 l culture medium: single strength (36.7 g for 1 l medium) for 6.8 l culture medium: double strength (73.4 g for 1 l medium) | 500 g |

isolation

| | | | |
|--------|---|--|-------|
| TN1130 | Endo Agar (Lactose Fuchsin Sulfite Agar) | for 14.5 l culture medium (34.6 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1259 | TBX Chromogen Agar (Tryptone Bile X Glucuronic Agar) | for 15.8 l culture medium (31.6 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1042 | Violet Red Bile Glucose Agar acc. to harm. EP/USP/JP (VRBD Agar) | for 13.7 l culture medium (36.5 g for 1 l culture medium) | 500 g |
| TN1149 | VRB Agar (Violet Red Bile Lactose Agar, VRBL Agar) | for 13.3 l culture medium (37.5 g for 1 l culture medium) | 500 g |
| TN1148 | VRBD Agar (Violet Red Bile Glucose Agar) | for 13.3 l culture medium (37.5 g for 1 l culture medium) | 500 g |

Salmonella

pre-enrichment

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1137 TN1137-01 | Buffered Peptone Water (BPW) | for 25.0 l culture medium (20.0 g for 1 l culture medium) for 250.0 l culture medium (20.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1013 | DIASSALM (Diagnostic Semi-Solid Salmonella Agar) | for 8.7 l culture medium (57.2 for 1 l culture medium) | 500 g |

enrichment

| Art. No. | Product | Description | Packing |
|----------|--|---|---------|
| TN1157 | Magnesium Chloride Malachite Green Broth acc. to Rappaport-Vassiliadis (RV Medium) | for 15.0 l culture medium (33.4 g for 1 l culture medium) | 500 g |
| TN1262 | Rappaport Vassiliadis Salmonella Enrichment Broth acc. to harm. EP/USP/JP (RVS Broth) | for 18.5 l culture medium (27.1 g for 1 l culture medium) | 500 g |
| TN1183 | Selenite Cystine Broth | for 21.7 l culture medium (23.0 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1187 | Tetrathionate Broth (Base) acc. to Muller-Kauffmann | for 5.0 l culture medium (89.4 g for 1 l culture medium) Addition of iodine-potassium iodide solution (5 g potassium iodide, 4 g iodine, 20 ml distilled water). | 500 g |

isolation

| | | | |
|---------------------|--|---|---------------|
| TN1110 | Brilliant Green Phenol Red Agar acc. to Edel Kampelmacher (BPLS Agar, mod.) | for 10.1 l culture medium (49.7 g for 1 l culture medium) | 500 g |
| TN1121 | Desoxycholate Citrate Agar Modified (Leifson Agar, DCLS Agar, mod.) | for 10.1 l culture medium (49.5 g for 1 l culture medium) | 500 g |
| TN1221 TN1221-01 | SS Agar (Salmonella Shigella Agar Modified) | for 8.8 l culture medium (56.5 g for 1 l culture medium) for 88.5 l culture medium (56.5 g for 1 l culture medium) | 500 g 5 kg |
| TN1170 | XLD Agar acc. to harm. EP/USP/JP (Xylose Lysine Deoxycholate Agar) | for 9.1 l culture medium (55.2 g for 1 l culture medium) | 500 g |
| TN1196 | XLD Agar acc. to ISO 6579 (Xylose Lysine Deoxycholate Agar) | for 8.8 l culture medium (57.0 g for 1 l culture medium) | 500 g |

confirmation

| | | | |
|--------|--|--|-------------|
| TN1146 | Kligler Iron Agar (KIA; Double Sugar Iron Agar acc. to Kligler) | for 10.1 l culture medium (49.6 g for 1 l culture medium) Urea Solution, 40% TN1308 may be added. | 500 g |
| TN1154 | Lysine Decarboxylase Broth (LDC Broth) | for 5.5 l culture medium (9.1 g for 1 l culture medium) | 50 g |
| TN1272 | MSRV Medium (Base) (Modified Semi-Solid Rappaport Vassiliadis Medium) | for 15.8 l culture medium (31.6 g for 1 l culture medium) Addition of Novobiocin Selective Supplement TN1324. | 500 g |
| TN1324 | Novobiocin Selective Supplement | for 6 l culture medium (2 vials for 1 l culture medium) 1 vial for 0.5 l MSRV Medium (Base) TN1272. | 12 x 1 vial |
| TN1308 | Urea Solution, 40 % | Supplement for culture medium: 1 vial for 1 l Kligler Iron Agar TN1146. | 12 x 1 vial |

Coagulase-positive staphylococci

enrichment

| Art. No. | Product | Description | Packing |
|-----------|---|--|---------|
| TN1083 | Tryptic Soy Broth Irradiated (CASO Broth, Soyabean Casein Digest Broth Irradiated) | for 16.7 l culture medium (30.0 g for 1 l culture medium) | 500 g |
| TN1083-01 | | for 166.7 l culture medium (30.0 g for 1 l culture medium) | 5 kg |

isolation

| | | | |
|--------|---|---|-----------|
| TN1104 | Baird Parker Agar (Base) | for 7.9 l culture medium (63.0 g for 1 l culture medium) Addition of Egg Yolk Tellurite Emulsion TN1310. | 500 g |
| TN1310 | Egg Yolk Potassium Tellurite Emulsion | for approx. 2 l culture medium (2 vials for 1 l culture medium) Supplement for culture media: 1 vial contains 25 ml 2 vials for 950 ml Baird Parker Agar (Base) TN1104. | 4 x 25 ml |
| TN1166 | Mannitol Salt Agar acc. to harm. EP/USP/JP | for 4.5 l culture medium (111.0 g for 1 l culture medium) | 500 g |

Pseudomonas aeruginosa

enrichment

| Art. No. | Product | Description | Packing |
|----------|------------------------------|---|---------|
| TN1158 | Malachite-green Broth | for 56.8 l culture medium (8.8 g for 1 l culture medium) No supplements necessary. | 500 g |

isolation

| | | | |
|--------|--|--|-------------|
| TN1424 | Glycerin, water-free Agar (Base) | Supplement for culture media: Pseudomonas Agar (Base) TN1286 | 100 ml |
| TN1517 | Oxidase reagent | for 12 tests 1 vial in 2 ml distilled water. | 12 x 1 vial |
| TN1286 | Pseudomonas Agar (Base) | for 10.3 l culture medium (48.4 g for 1 l culture medium) Addition of Pseudomonas CN Selective Supplement TN1323 and glycerin TN1424. | 500 g |
| TN1323 | Pseudomonas CN Selective Supplement | for 12 tests (2 vials for 1 l culture medium) 1 vial in 2 ml distilled water. | 12 x 1 vial |

Lactobacilli

| Art. No. | Product | Description | Packing |
|----------|---|---|---------|
| TN1201 | MRS Agar (Base) (Lactobacillus Agar acc. to de Man, Rogosa and Sharpe) | for 8.0 l culture medium (62.3 g for 1 l culture medium) Addition of Tween® 80 TN1422. | 500 g |
| TN1068 | MRS Agar (Base) with pH 5.7 | for 7.8 l culture medium (64.3 g for 1 l culture medium) Addition of Tween® 80 TN1422. | 500 g |
| TN1205 | MRS Broth (Base) | for 9.7 l culture medium (51.3 g for 1 l culture medium) Addition of Tween® 80 TN1422. | 500 g |
| TN1422 | Tween® 80 | 1 ml for 1 l MRS Agar (Base) TN1201. 1 ml for 1 l MRS Agar (Base) with pH 5.7 TN1068. 1 ml for 1 l MRS Broth (Base) TN1205. | 100 ml |

Yeasts and moulds

| Art. No. | Product | Description | Packing |
|---------------------|--|---|---------------|
| TN1424 | Glycerin, water-free | Supplement for culture media: Wort Agar (Base) TN1210 | 100 ml |
| TN1264 | Sabouraud Dextrose Agar acc. to harm. EP/USP/JP | for 7.7 l culture medium (65.0 g for 1 l culture medium) | 500 g |
| TN1178 TN1178-01 | Sabouraud 2% Glucose Agar (Sabouraud Dextrose Agar) | for 11.9 l culture medium (42.0 g for 1 l culture medium) for 119.0 l culture medium (42.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1746 | Wort Agar | 1 bottle contains 250 ml. | 250 ml |
| TN1210 | Wort Agar (Base) | for 10.4 l culture medium (48.0 g for 1 l culture medium) Addition of glycerin TN1424. | 500 g |
| TN1144 | Yeast Extract Glucose Chloramphenicol Agar (YGC Agar) | for 14.2 l culture medium (35.1 g for 1 l culture medium) No supplements necessary. | 500 g |

Clostrides

| Art. No. | Product | Description | Packing |
|----------|---|---|---------|
| TN1116 | Clostridial Differential Broth (DRCM) | for 16.6 l culture medium: single strength (30.1 g for 1 l medium) for 8.3 l culture medium: double strength (60.2 g for 1 l medium) | 500 g |
| TN1167 | Reinforced Medium for Clostridia acc. to harm. EP/USP/JP (RCM) | for 13.2 l culture medium (38.0 g for 1 l culture medium) | 500 g |

Mesophile spore formers

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1031 TN1031-01 | Casein Soya Bean Digest Agar acc. to harm. EP/USP/JP (CASO Agar) | for 12.5 l culture medium (40.0 g for 1 l culture medium) for 125.0 l culture medium (40.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1763 | Casein Soya Bean Digest Broth with 6.5 % NaCl | 1 bottle contains 2 ml broth. | 100 bottles |

Examination of Meat and Meat Products

Germ number determination

| Art. No. | Product | Description | Packing |
|---------------------|---|--|---------------|
| TN1164 TN1164-01 | Nutrient Agar I | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) Blood may be added. | 500 g 5 kg |
| TN1189 | Plate Count Agar (TGE Agar, Tryptone Glucose Yeast Extract Agar) | for 27.0 l culture medium (18.5 g for 1 l culture medium) | 500 g |

Enterobacteriaceae

pre-enrichment

| Art. No. | Product | Description | Packing |
|---------------------|-------------------------------------|---|---------------|
| TN1137 TN1137-01 | Buffered Peptone Water (BPW) | for 25.0 l culture medium (20.0 g for 1 l culture medium) for 250.0 l culture medium (20.0 g for 1 l culture medium) | 500 g 5 kg |

isolation

| | | | |
|---------------------|--|---|---------------|
| TN1086 | Glucose Agar | for 13.0 l culture medium (38.5 g for 1 l culture medium) | 500 g |
| TN1156 | MacConkey Agar | for 10.3 l culture medium (48.5 g for 1 l culture medium) | 500 g |
| TN1075 TN1075-01 | MacConkey Agar acc. to harm. EP/USP/JP | for 10.0 l culture medium (50.0 g for 1 l culture medium) for 100.0 l culture medium (50.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1042 | Violet Red Bile Glucose Agar acc. to harm. EP/USP/JP (VRBD) | for 13.7 l culture medium (36.5 g for 1 l culture medium) | 500 g |
| TN1148 | VRBD Agar (Violet Red Bile Glucose Agar) | for 13.3 l culture medium (37.5 g for 1 l culture medium) | 500 g |
| TN1194 TN1194-01 | Water-blue Metachrome-yellow Lactose Agar acc. to Gassner Mod. (Gassner Agar, mod.) | for 9.9 l culture medium (50.3 g for 1 l culture medium) for 99.4 l culture medium (50.3 g for 1 l culture medium) | 500 g 5 kg |

confirmation

| | | | |
|--------|--|--|-------|
| TN1146 | Kligler Iron Agar (KIA, Double Sugar Iron Agar acc. to Kligler) | for 10.1 l culture medium (49.6 g for 1 l culture medium) Urea TN1308 may be added. | 500 g |
| TN1184 | Simmons Citrate Agar | for 9.8 l culture medium (15.3 g for 1 l culture medium) | 150 g |

Salmonella

pre-enrichment

| Art. No. | Product | Description | Packing |
|---------------------|-------------------------------------|---|---------------|
| TN1137 TN1137-01 | Buffered Peptone Water (BPW) | for 25.0 l culture medium (20.0 g for 1 l culture medium) for 250.0 l culture medium (20.0 g for 1 l culture medium) | 500 g 5 kg |

enrichment

| | | | |
|--------|--|---|-------|
| TN1157 | Magnesium Chloride Malachite Green Broth acc. to Rappaport Vassiliadis (RV Medium) | for 15.0 l culture medium (33.4 g for 1 l culture medium) | 500 g |
| TN1262 | Rappaport Vassiliadis Salmonella Enrichment Broth acc. to harm. EP/USP/JP (RVS Broth) | for 18.5 l culture medium (27.1 g for 1 l culture medium) | 500 g |
| TN1183 | Selenite Cystine Broth | for 21.7 l culture medium (23.0 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1187 | Tetrathionate Broth (Base) acc. to Muller-Kauffmann | for 5.0 l culture medium (89.4 g for 1 l culture medium) Addition of iodine-potassium iodide solution (5 g potassium iodide, 4 g iodine, 20 ml distilled water). | 500 g |

isolation

| | | | |
|---------------------|---|---|---------------|
| TN1111 | Brilliant Green Phenol Red Agar (BPLS Agar acc. to Kauffmann, mod.) | for 10.0 l culture medium (50.0 g for 1 l culture medium) | 500 g |
| TN1110 | Brilliant Green Phenol Red Agar acc. to Edel Kampelmacher (BPLS Agar, mod.) | for 10.1 l culture medium (49.7 g for 1 l culture medium) | 500 g |
| TN1121 | Desoxycholate Citrate Agar Modified (Leifson Agar, DCLS Agar, mod.) | for 10.1 l culture medium (49.5 g for 1 l culture medium) | 500 g |
| TN1221 TN1221-01 | SS Agar (Salmonella Shigella Agar Modified) | for 8.8 l culture medium (56.5 g for 1 l culture medium) for 88.5 l culture medium (56.5 g for 1 l culture medium) | 500 g 5 kg |
| TN1194 TN1194-01 | Water-blue Metachrome-yellow Lactose Agar acc. to Gassner, Modified (Gassner Agar, mod.) | for 9.9 l culture medium (50.3 g for 1 l culture medium) for 99.4 l culture medium (50.3 g for 1 l culture medium) | 500 g 5 kg |
| TN1170 | XLD Agar acc. to harm. EP/USP/JP (Xylose Lysine Deoxycholate Agar) | for 9.1 l culture medium (55.2 g for 1 l culture medium) | 500 g |
| TN1196 | XLD Agar acc. to ISO 6579 (Xylose Lysine Deoxycholate Agar) | for 8.8 l culture medium (57.0 g for 1 l culture medium) | 500 g |

confirmation

| Art. No. | Product | Description | Packing |
|----------|---|--|-------------|
| TN1146 | Kligler Iron Agar (KIA; Double Sugar Iron Agar acc. to Kligler) | for 10.1 l culture medium (49.6 g for 1 l culture medium) Urea TN1308 may be added. | 500 g |
| TN1154 | Lysine Decarboxylase Broth (LDC Broth) | for 5.5 l culture medium (9.0 g for 1 l culture medium) | 50 g |
| TN1272 | MSRV Medium (Base) (Modified Semi-Solid Rappaport Vassiliadis Medium Base) | for 15.8 l culture medium (31.7 g for 1 l culture medium) Addition of Novobiocin Selective Supplement TN1324. | 500 g |
| TN1324 | Novobiocine Selective Supplement | for 6 l culture medium (2 vials for 1 l culture medium) 1 vial for 0.5 l MSRV Medium (Base) TN1272. | 12 x 1 vial |
| TN1308 | Urea Solution, 40 % | Supplement for culture medium: 1 vial for 1 l Kligler Iron Agar TN1146. | 12 x 1 vial |

Coagulase-positive staphylococci

enrichment

| Art. No. | Product | Description | Packing |
|-----------|---|--|---------|
| TN1083 | Tryptic Soy Broth Irradiated (CASO Broth, Soyabean Casein Digest Broth Irradiated) | for 16.7 l culture medium (30.0 g for 1 l culture medium) | 500 g |
| TN1083-01 | | for 166.7 l culture medium (30.0 g for 1 l culture medium) | 5 kg |

isolation

| | | | |
|--------|---|---|-----------|
| TN1104 | Baird Parker Agar (Base) | for 7.9 l culture medium (63.0 g for 1 l culture medium) Addition of Egg Yolk Tellurite Emulsion TN1310. | 500 g |
| TN1310 | Egg Yolk Potassium Tellurite Emulsion | for approx. 2 l culture medium (2 vials for 1 l culture medium) Supplement for culture media: 1 vial contains 25 ml 2 vials for 950 ml Baird Parker Agar (Base) TN1104. | 4 x 25 ml |
| TN1166 | Mannitol Salt Agar acc. to harm. EP/USP/JP | for 4.5 l culture medium (111.0 g for 1 l culture medium) | 500 g |

Listeria

| Art. No. | Product | Description | Packing |
|---------------------|--|--|---------------|
| TN1257 | Fraser Broth (Base) | for 8.7 l culture medium (57.4 g for 1 l culture medium) Addition of Fraser Selective Supplement TN1318. | 500 g |
| TN1035 | Fraser Broth Complete | for 8.6 l culture medium (57.9 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1034 TN1034-01 | ½ Fraser Broth Complete | for 8.6 l culture medium (57.9 g for 1 l culture medium) for 86.3 l culture medium (57.9 g for 1 l culture medium) No supplements necessary. | 500 g 5 kg |
| TN1318 | Fraser Selective Supplement | for 3 l culture medium (1 vial per component for 0.5 l medium) Supplement for culture media consists of two components. 1 vial per component for 0.5 l Fraser Broth (Base) TN1257. | 6 x 2 vials |
| TN1209 | Palcam Agar (Base) (Palcam Agar acc. to van Netten) | for 7.4 l culture medium (68.0 g for 1 l culture medium) Addition of Palcam Selective Supplement TN1312. | 500 g |
| TN1312 | Palcam Selective Supplement | for 6 l culture medium (2 vials for 1 l culture medium) 1 vial for 0.5 l Palcam Agar (Base) TN1209. | 12 x 1 vial |

Bacillus cereus

| Art. No. | Product | Description | Packing |
|----------|--|--|-------------|
| TN1245 | Bacillus Cereus Agar (Base) (PEMBA) | for 12.5 l culture medium (39.9 g for 1 l culture medium) Addition of Egg Yolk Emulsion TN1316 and Bacillus Cereus Selective Supplement TN1315. | 500 g |
| TN1315 | Bacillus Cereus Selective Supplement (Polymyxin B Selective Supplement) | for 6 l culture medium (2 vials for 1 culture medium) In conjunction with Bacillus Cereus Agar (Base) TN1245 and Egg Yolk Emulsion TN1316. | 12 x 1 vial |
| TN1316 | Egg Yolk Emulsion | for approx. 1-2 l culture medium (2-4 vials for 1 l culture medium) Supplement for culture media: 1 vial contains 25 ml 2 vials for 950 ml Bacillus Cereus Agar (Base) TN1245. 4 vials for 900 ml MYP Agar (Base) TN1291. | 4 x 25 ml |
| TN1291 | MYP Agar (Base) (Cereus Selective Agar Base acc. to Mossel) | for 11.6 l culture medium (43.0 g for 1 l culture medium) Addition of Egg Yolk Emulsion TN1316 and Bacillus Cereus Selective Supplement TN1315. | 500 g |

For Use in the Beverage Industry

| Art. No. | Product | Description | Packing |
|---------------------|---|---|-----------------------|
| TN1724 | DEV Nutrient Agar | 1 bottle contains 250 ml. | 250 ml |
| TN1128 TN1128-01 | DEV Nutrient Agar | for 14.3 l culture medium (35.0 g for 1 l culture medium) for 142.9 l culture medium (35.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1424 | Glycerin, water-free | Supplement for culture media: Wort Agar (Base) TN1210 | 100 ml |
| TN1066 | Linden Grain Medium | for 847.5 l culture medium (29.5 g for 1 l culture medium) | 25 kg |
| TN1743 | Lysine Agar | 1 bottle contains 250 ml. | 250 ml |
| TN1707 TN1707-01 | MegaPec Broth | 1 tube contains 12 ml broth. 1 tube contains 12 ml broth. | 20 tubes 100 tubes |
| TN1201 | MRS Agar (Base) (Lactobacillus Agar acc. to de Man, Rogosa and Sharpe) | for 8.0 l culture medium (62.3 g for 1 l culture medium) Addition of Tween® 80 TN1422. | 500 g |
| TN1207 | Orange Serum Agar | for 12.5 l culture medium (40.0 g for 1 l culture medium) | 500 g |
| TN1269 | Orange Serum Broth | for 20.0 l culture medium (25.0 g for 1 l culture medium) | 500 g |
| TN1502 | VLB-S7-S Agar | 1 bottle contains 250 ml. | 250 ml |
| TN1503 | VLB-S7-S Broth | 1 bottle contains 250 ml. | 250 ml |
| TN1744 | WL Nutrient Agar (WLN) | 1 bottle contains 250 ml. | 250 ml |
| TN1745 | WL Nutrient Agar with Cycloheximide (WLD) | 1 bottle contains 250 ml. | 250 ml |
| TN1746 | Wort Agar | 1 bottle contains 250 ml. | 250 ml |
| TN1210 | Wort Agar (Base) | for 10.4 l culture medium (48.0 g for 1 l culture medium) Addition of glycerin TN1424. | 500 g |



Culture Media for Use in the Beverage Industry

Microbiology

Ready-to-use culture media

Dehydrated culture media

More safety in the beverage industry

- Very well prepared for quality control in the beverage industry
- Reliable results in the microbiological examination of non-alcoholic or alcoholic beverages
- Culture media in order to specifically support the growth of particularly fastidious species
- Ready-to-use culture media in tubes and bottles or as dehydrated culture media available

Do you have a specific request?

We develop and produce according to your demand.



Detection of Anaerobic Microorganisms

| Art. No. | Product | Description | Packing |
|---------------------|--|---|---------------|
| TN1031 TN1031-01 | Casein Soya Bean Digest Agar acc. to harm. EP/USP/JP (CASO Agar) | for 12.5 l culture medium (40.0 g for 1 l culture medium) for 125.0 l culture medium (40.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1116 | Clostridial Differential Broth (DRCM) | for 16.6 l culture medium: single strength (30.1 g for 1 l medium) for 8.3 l culture medium: double strength (60.2 g for 1 l medium) | 500 g |
| TN1139 | Glucose Yeast Extract Cysteine Agar (Base) acc. to Beerens | for 13.8 l culture medium (36.3 g for 1 l culture medium) Blood addition necessary. | 500 g |
| TN1167 | Reinforced Medium for Clostridia acc. to harm. EP/USP/JP (RCM) | for 13.2 l culture medium (38.0 g for 1 l culture medium) | 500 g |
| TN1288 | m-CP Agar (Base) (Chromogenic Membrane Filtration Clostridium perfringens Agar) | for 7.3 l culture medium (68.2 g for 1 l culture medium) Addition of m-CP-Selective Supplement TN1330. | 500 g |
| TN1330 | m-CP Selective Supplement | for 2 l culture medium (1 vial per component for 0.5 l culture medium) Supplement for culture media consists of three components. 1 vial per component for 0.5 l m-CP Agar (Base) TN1288. | 4 x 3 vials |
| TN1203 | Schaedler Broth | for 19.1 l culture medium (26.2 g for 1 l culture medium) | 500 g |
| TN1188 TN1188-01 | Thioglycollate Medium acc. to EP/USP (Fluid Thioglycollate Medium) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1256 | Thioglycollate Medium Modified (high degree of transparency) | for 16.7 l culture medium (30.0 g for 1 l culture medium) | 500 g |
| TN1023 | Thioglycollate Medium without Indicator | for 16.6 l culture medium (30.1 g for 1 l culture medium) | 500 g |
| TN1241 | TSC Agar (Base) (Tryptose Sulfit Cycloserine Agar) | for 12.8 l culture medium (39.0 g for 1 l culture medium) Addition of TSC Selective Supplement TN1333. | 500 g |
| TN1333 | TSC Selective Supplement | for 6 l culture medium (2 vials for 1 l culture medium) 1 vial for 0.5 l TSC Agar (Base) TN1241. | 12 x 1 vial |

Detection of Yeasts and Moulds

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1263 TN1263-01 | Casein Soya Bean Digest Broth acc. to harm. EP/USP/JP (CASO Broth) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1424 | Glycerin, water-free | Supplement for culture media: Wort Agar (Base) TN1210 | 100 ml |
| TN1150 | Potato Dextrose Agar acc. to harm. EP/USP/JP | for 12.8 l culture medium (39.0 g for 1 l culture medium) | 500 g |
| TN1264 | Sabouraud Dextrose Agar acc. to harm. EP/USP/JP | for 7.7 l culture medium (65.0 g for 1 l culture medium) | 500 g |
| TN1047 | Sabouraud Dextrose Broth acc. to harm. EP/USP/JP | for 16.7 l culture medium (30.0 g for 1 l culture medium) | 500 g |
| TN1178 TN1178-01 | Sabouraud 2% Glucose Agar (Sabouraud Dextrose Agar) | for 11.9 l culture medium (42.0 g for 1 l culture medium) for 119.0 l culture medium (42.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1015 | Sabouraud Glucose Medium with Chloramphenicol | for 8.1 l culture medium (62.1 g for 1 l culture medium) 405 mg Chloramphenicol for 8.1 l culture medium | 500 g |
| TN1210 | Wort Agar (Base) | for 10.4 l culture medium (48.0 g for 1 l culture medium) Addition of glycerin TN1424. | 500 g |
| TN1144 | YGC Agar (YGC Agar) | for 14.2 l culture medium (35.1 g for 1 l culture medium) No supplements necessary. | 500 g |

Biochemical Identification

| Art. No. | Product | Description | Packing |
|----------|--|--|-------------|
| TN1113 | Bromthymol Blue Broth (Base) | for 33.3 l culture medium (15.0 g for 1 l culture medium) | 500 g |
| TN1146 | Kligler Iron Agar (KIA, Double Sugar Iron Agar acc. to Kligler) | for 10.1 l culture medium (49.6 g for 1 l culture medium) Urea Solution, 40% TN1308 may be added. | 500 g |
| TN1154 | Lysine Decarboxylase Broth (LDC Broth) | for 5.5 l culture medium (9.1 g for 1 l culture medium) | 50 g |
| TN1184 | Simmons Citrate Agar | for 9.8 l culture medium (15.3 g for 1 l culture medium) | 150 g |
| TN1308 | Urea Solution, 40 % | Supplement for culture medium: 1 vial for 1 l Kligler Iron Agar TN1146. | 12 x 1 vial |

Culture Media for Pharma

MICROBIOLOGY

You don't need a prescription to obtain our culture media

Contents

- *Escherichia coli*
- *Salmonella*
- *Pseudomonas aeruginosa*
- *Staphylococcus aureus*
- Clostrides
- *Candida albicans*
- Enumeration
- Bile-tolerant gram-negative bacteria
- Dermatophytes



Escherichia coli

pre-enrichment

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1263 TN1263-01 | Casein Soya Bean Digest Broth acc. to harm. EP/USP/JP (CASO Broth) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |

isolation

| | | | |
|---------------------|--|---|---------------|
| TN1075 TN1075-01 | MacConkey Agar acc. to harm. EP/USP/JP | for 10.0 l culture medium (50.0 g for 1 l culture medium) for 100.0 l culture medium (50.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1027 | MacConkey Broth acc. to harm. EP/USP/JP | for 14.3 l culture medium (35.0 g for 1 l culture medium) | 500 g |

Salmonella

pre-enrichment

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1263 TN1263-01 | Casein Soya Bean Digest Broth acc. to harm. EP/USP/JP (CASO Broth) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |

selective enrichment

| | | | |
|--------|--|---|-------|
| TN1262 | Rappaport Vassiliadis Salmonella Enrichment Broth acc. to harm. EP/USP/JP (RVS Broth) | for 18.5 l culture medium (27.1 g for 1 l culture medium) | 500 g |
|--------|--|---|-------|

isolation

| | | | |
|--------|---|--|-------|
| TN1170 | XLD Agar acc. to harm. EP/USP/JP (Xylose Lysine Deoxycholate Agar) | for 9.1 l culture medium (55.2 g for 1 l culture medium) | 500 g |
|--------|---|--|-------|

Pseudomonas aeruginosa

pre-enrichment

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1263 TN1263-01 | Casein Soya Bean Digest Broth acc. to harm. EP/USP/JP (CASO Broth) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |

Staphylococcus aureus

pre-enrichment

| Art. No. | Product | Description | Packing |
|-----------|---|--|---------|
| TN1263 | Casein Soya Bean Digest Broth | for 16.7 l culture medium (30.0 g for 1 l culture medium) | 500 g |
| TN1263-01 | acc. to harm. EP/USP/JP (CASO Broth) | for 166.7 l culture medium (30.0 g for 1 l culture medium) | 5 kg |

selective enrichment

| | | | |
|--------|---|---|-------|
| TN1166 | Mannitol Salt Agar acc. to harm. EP/USP/JP | for 4.5 l culture medium (111.0 g for 1 l culture medium) | 500 g |
|--------|---|---|-------|

Clostrides

selective enrichment

| Art. No. | Product | Description | Packing |
|----------|---|---|---------|
| TN1167 | Reinforced Medium for Clostridia | for 13.2 l culture medium (38.0 g for 1 l culture medium) | 500 g |
| | acc. to harm. EP/USP/JP | Blood may be added. | |

isolation

| | | | |
|--------|--|---|-------|
| TN1118 | Columbia Agar (Base) | for 12.5 l culture medium (40.0 g for 1 l culture medium) | 500 g |
| | (Columbia Blood Agar Base) | Addition of blood. | |
| TN1191 | Columbia Agar acc. to harm. EP/USP/JP | for 12.2 l culture medium (41.0 g for 1 l culture medium) | 500 g |

Candida albicans

pre-enrichment

| Art. No. | Product | Description | Packing |
|----------|---------------------------------|---|---------|
| TN1047 | Sabouraud Dextrose Broth | for 16.7 l culture medium (30.0 g for 1 l culture medium) | 500 g |
| | acc. to harm. EP/USP/JP | | |

selective enrichment

| | | | |
|-----------|----------------------------------|--|-------|
| TN1264 | Sabouraud Dextrose Agar | for 7.7 l culture medium (65.0 g for 1 l culture medium) | 500 g |
| | acc. to harm. EP/USP/JP | | |
| TN1178 | Sabouraud 2% Glucose Agar | for 11.9 l culture medium (42.0 g for 1 l culture medium) | 500 g |
| TN1178-01 | (Sabouraud Dextrose Agar) | for 119.0 l culture medium (42.0 g for 1 l culture medium) | 5 kg |

Enumeration

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1031 TN1031-01 | Casein Soya Bean Digest Agar acc. to harm. EP/USP/JP (CASO Agar) | for 12.5 l culture medium (40.0 g for 1 l culture medium) for 125.0 l culture medium (40.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1263 TN1263-01 | Casein Soya Bean Digest Broth acc. to harm. EP/USP/JP (CASO Broth) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1150 | Potato Dextrose Agar acc. to harm. EP/USP/JP | for 12.8 l culture medium (39.0 g for 1 l culture medium) | 500 g |
| TN1025 TN1025-01 | Buffered Sodium Chloride Peptone Solution pH 7.0 acc. to harm. EP/USP/JP | for 34.2 l culture medium (14.6 g for 1 l culture medium) for 342.4 l culture medium (14.6 g for 1 l culture medium) | 500 g 5 kg |
| TN1264 | Sabouraud Dextrose Agar acc. to harm. EP/USP/JP | for 7.7 l culture medium (65.0 g for 1 l culture medium) | 500 g |

Bile-tolerant gram-negative bacteria

pre-enrichment

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1263 TN1263-01 | Casein Soya Bean Digest Broth acc. to harm. EP/USP/JP (CASO Broth) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |

enrichment

| | | | |
|--------|---|---|-------|
| TN1236 | Enterobacteriaceae Enrichment Broth-Mossel acc. to harm. EP/USP/JP | for 11.5 l culture medium (43.4 g for 1 l culture medium) | 500 g |
|--------|---|---|-------|

isolation

| | | | |
|--------|--|---|-------|
| TN1042 | Violet Red Bile Glucose Agar acc. to harm. EP/USP/JP (VRBD) | for 13.7 l culture medium (36.5 g for 1 l culture medium) | 500 g |
|--------|--|---|-------|

Dermatophytes

| Art. No. | Product | Description | Packing |
|---------------------|--|---|---------------|
| TN2102 | Dermatophytes Agar, modified | Culture medium with Chloramphenicol and Cycloheximide included. No supplements necessary. | 2 x 10 plates |
| TN1178 TN1178-01 | Sabouraud 2% Glucose Agar (Sabouraud Dextrose Agar) | for 11.9 l culture medium (42.0 g for 1 l culture medium) for 119.0 l culture medium (42.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1016 | Sabouraud 2% Glucose Chloramphenicol Agar | for 11.9 l culture medium (42.1 g for 1 l culture medium) 1190 mg Chloramphenicol for 11.9 l culture medium | 500 g |

Culture Media for Cosmetics

MICROBIOLOGY



... but true beauty comes from within

Contents

- *Escherichia coli*
- Mesophile spore formers
- *Pseudomonas aeruginosa*
- *Staphylococcus aureus*



E. coli

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1031 TN1031-01 | Casein Soya Bean Digest Agar acc. to harm. EP/USP/JP (CASO Agar) | for 12.5 l culture medium (40.0 g for 1 l culture medium) for 125.0 l culture medium (40.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1156 | MacConkey Agar | for 10.3 l culture medium (48.5 g for 1 l culture medium) | 500 g |
| TN1075 TN1075-01 | MacConkey Agar acc. to harm. EP/USP/JP | for 10.0 l culture medium (50.0 g for 1 l culture medium) for 100.0 l culture medium (50.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1147 | Peptone Saline Diluent (Maximum Recovery Diluent) | for 52.6 l culture medium (9.5 g for 1 l culture medium) | 500 g |

Mesophile spore formers

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1031 TN1031-01 | Casein Soya Bean Digest Agar acc. to harm. EP/USP/JP (CASO Agar) | for 12.5 l culture medium (40.0 g for 1 l culture medium) for 125.0 l culture medium (40.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1025 TN1025-01 | Buffered Sodium Chloride Peptone Solution pH 7.0 acc. to harm. EP/USP/JP | for 34.2 l culture medium (14.6 g for 1 l culture medium) for 342.4 l culture medium (14.6 g for 1 l culture medium) | 500 g 5 kg |
| TN1147 | Peptone Saline Diluent (Maximum Recovery Diluent) | for 52.6 l culture medium (9.5 g for 1 l culture medium) | 500 g |

Pseudomonas aeruginosa

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1031 TN1031-01 | Casein Soya Bean Digest Agar acc. to harm. EP/USP/JP (CASO Agar) | for 14.3 l culture medium (40.0 g for 1 l culture medium) for 142.9 l culture medium (40.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1147 | Peptone Saline Diluent (Maximum Recovery Diluent) | for 52.6 l culture medium (9.5 g for 1 l culture medium) | 500 g |

Staphylococcus aureus

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1104 | Baird Parker Agar (Base) | for 7.9 l culture medium (63.0 g for 1 l culture medium) Addition of Egg Yolk Tellurite Emulsion TN1310. | 500 g |
| TN1031 TN1031-01 | Casein Soya Bean Digest Agar acc. to harm. EP/USP/JP (CASO Agar) | for 14.3 l culture medium (40.0 g for 1 l culture medium) for 142.9 l culture medium (40.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1310 | Egg Yolk Potassium Tellurite Emulsion | for approx. 2 l culture medium (2 vials for 1 l culture medium) Supplement for culture media: 1 vial contains 25 ml 2 vials for 950 ml Baird Parker Agar (Base) TN1104. | 4 x 25 ml |
| TN1166 | Mannitol Salt Agar acc. to harm. EP/USP/JP | for 4.5 l culture medium (111.0 g for 1 l culture medium) | 500 g |
| TN1147 | Peptone Saline Diluent (Maximum Recovery Diluent) | for 52.6 l culture medium (9.5 g for 1 l culture medium) | 500 g |

Culture Media in Veterinary

MICROBIOLOGY



We would be glad to help you count

Contents

- Enterobacteriaceae
- Moulds
- Universal culture media
- Pseudomonads
- *Salmonella* and *Shigella*
- Staphylococci
- Streptococci



Enterobacteriaceae

| Art. No. | Product | Description | Packing |
|---------------------|--|---|---------------|
| TN1110 | Brilliant Green Phenol Red Agar acc. to Edel Kampelmacher (BPLS Agar, mod.) | for 10.1 l culture medium (49.7 g for 1 l culture medium) | 500 g |
| TN1121 | Desoxycholate Citrate Agar Modified (Leifson Agar, DCLS Agar, mod.) | for 10.1 l culture medium (49.5 g for 1 l culture medium) | 500 g |
| TN1130 | Endo Agar (Lactose Fuchsin Sulfite Agar) | for 14.5 l culture medium (34.6 g for 1 l culture medium) No supplements necessary. | 500 g |
| TN1157 | Magnesium Chloride Malachite Green Broth acc. to Rappaport-Vassiliadis (RV Medium) | for 15.0 l culture medium (33.4 g for 1 l culture medium) | 500 g |
| TN1040 | Minca Agar, modified | for 18.9 l culture medium (26.4 g for 1 l culture medium) Addition of Minca Supplement TN 1334. | 500 g |
| TN1334 | Minca Supplement | for 1.2 l culture medium (1 vial for 0.1 l culture medium) 1 vial for 100 ml Minca Agar, modified TN1040. | 12 x 1 vial |
| TN1262 | Rappaport Vassiliadis Salmonella Enrichment Broth acc. to harm. EP/USP/JP (RVS Broth) | for 18.5 l culture medium (27.1 g for 1 l culture medium) | 500 g |
| TN1221 TN1221-01 | SS Agar (Salmonella Shigella Agar Modified) | for 8.8 l culture medium (56.5 g for 1 l culture medium) for 88.5 l culture medium (56.5 g for 1 l culture medium) | 500 g 5 kg |
| TN1042 | Violet Red Bile Glucose Agar acc. to harm. EP/USP/JP (VRBD Agar) | for 13.7 l culture medium (36.5 g for 1 l culture medium) | 500 g |
| TN1148 | VRBD Agar (Violet Red Bile Glucose Agar) | for 13.3 l culture medium (37.5 g for 1 l culture medium) | 500 g |
| TN1194 TN1194-01 | Water-blue Metachrome-yellow Lactose Agar acc. to Gassner Mod. (Gassner Agar, mod.) | for 9.9 l culture medium (50.3 g for 1 l culture medium) for 99.4 l culture medium (50.3 g for 1 l culture medium) | 500 g 5 kg |

Moulds

| Art. No. | Product | Description | Packing |
|---------------------|--|---|---------------|
| TN1424 | Glycerin, water-free | Supplement for culture media: Wort Agar (Base) TN1210 | 100 ml |
| TN1264 | Sabouraud Dextrose Agar acc. to harm. EP/USP/JP | for 7.7 l culture medium (65.0 g for 1 l culture medium) | 500 g |
| TN1178 TN1178-01 | Sabouraud 2% Glucose Agar (Sabouraud Dextrose Agar) | for 11.9 l culture medium (42.0 g for 1 l culture medium) for 119.0 l culture medium (42.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1143 | Urea Agar (Base) acc. to Christensen | for 14.7 l culture medium (17.0 g for 1 l culture medium) Addition of Urea Solution, 40% TN1308. | 250 g |
| TN1308 | Urea Solution, 40 % | Supplement for culture medium. | 12 x 1 vial |
| TN1746 | Wort Agar | 1 bottle contains 250 ml. | 250 ml |
| TN1210 | Wort Agar (Base) | for 10.4 l culture medium (48.0 g for 1 l culture medium) Addition of glycerin TN1424. | 500 g |

Universal culture media

| Art. No. | Product | Description | Packing |
|---------------------|---|---|---------------|
| TN1216 | Brain Heart Infusion Broth (BHI Broth, Brain Heart Glucose Broth) | for 13.5 l culture medium (37.0 g for 1 l medium) | 500 g |
| TN1188 TN1188-01 | Thioglycollate Medium acc. to EP/USP (Fluid Thioglycollate Medium) | for 16.7 l culture medium (30.0 g for 1 l culture medium) for 166.7 l culture medium (30.0 g for 1 l culture medium) | 500 g 5 kg |
| TN1256 | Thioglycollate Medium Modified (high degree of transparency) | for 16.7 l culture medium (30.0 g for 1 l culture medium) | 500 g |
| TN1023 | Thioglycollate Medium without Indicator | for 16.6 l culture medium (30.1 g for 1 l culture medium) | 500 g |

Pseudomonads

| Art. No. | Product | Description | Packing |
|----------|--|---|-------------|
| TN1424 | Glycerin, water-free | Supplement for culture media: Pseudomonas Agar (Base) TN1286 | 100 ml |
| TN1286 | Pseudomonas Agar (Base) | for 10.3 l culture medium (48.4 g for 1 l culture medium) Addition of Pseudomonas CN Selective Supplement TN1323 and glycerin TN1424. | 500 g |
| TN1323 | Pseudomonas CN Selective Supplement | for 6 l culture medium (2 vials for 1 l culture medium) 1 vial for 0.5 l Pseudomonas Agar (Base) TN1286. | 12 x 1 vial |

Salmonella and Shigella

| Art. No. | Product | Description | Packing |
|----------|---|--|---------|
| TN1170 | XLD Agar acc. to harm. EP/USP/JP (Xylose Lysine Deoxycholate Agar) | for 9.1 l culture medium (55.2 g for 1 l culture medium) | 500 g |
| TN1196 | XLD Agar acc. to ISO 6579 (Xylose Lysine Deoxycholate Agar) | for 8.8 l culture medium (57.0 g for 1 l culture medium) | 500 g |

Staphylococci

| Art. No. | Product | Description | Packing |
|----------|---|---|-----------|
| TN1104 | Baird Parker Agar (Base) | for 7.9 l culture medium (63.0 g for 1 l culture medium) Addition of Egg Yolk Tellurite Emulsion TN1310. | 500 g |
| TN1310 | Egg Yolk Potassium Tellurite Emulsion | for approx. 2 l culture medium (2 vials for 1 l culture medium) Supplement for culture media: 1 vial contains 25 ml 2 vials for 950 ml Baird Parker Agar (Base) TN1104. | 4 x 25 ml |
| TN1166 | Mannitol Salt Agar acc. to harm. EP/USP/JP | for 4.5 l culture medium (111.0 g for 1 l culture medium) | 500 g |

Streptococci

| Art. No. | Product | Description | Packing |
|----------|--|---|---------|
| TN1118 | Columbia Agar (Base) (Columbia Blood Agar Base) | for 12.5 l culture medium (40.0 g for 1 l culture medium) Addition of blood. | 500 g |
| TN1191 | Columbia Agar acc. to harm. EP/USP/JP | for 12.2 l culture medium (41.0 g for 1 l culture medium) | 500 g |

Certificate

Certificate DIN EN ISO/IEC 17025:
<http://tritium-microbiologie.nl/en/>

Notes

Do you have a specific request?
We develop and produce according to your demand.

sifin

sifin diagnostics gmbh

Berliner Allee 317-321
13088 Berlin, Germany

Phone: +49 30 700 144-0
Telefax: +49 30 700 144-30
E-Mail: info@sifin.de



sifin diagnostics gmbh

Berliner Allee 317-321
13088 Berlin, Germany

Phone: +49 30 700 144-0
Telefax: +49 30 700 144-30
E-Mail: info@sifin.de

www.sifin.de